

Evidence-Informed Milestones for Developmental Surveillance Tools

Jennifer M. Zubler, MD,^{a,b} Lisa D. Wiggins, PhD,^a Michelle M. Macias, MD,^{c,*} Toni M. Whitaker, MD,^d Judith S. Shaw, EdD, MPH, RN,^e Jane K. Squires, PhD,^f Julie A. Pajek, PhD,^g Rebecca B. Wolf, MA,^k Karnesha S. Slaughter, MPH,^a Amber S. Broughton, MPH,^a Krysta L. Gerndt, MPH,ⁱ Bethany J. Mlodoch,^h Paul H. Lipkin, MD^{l,j,*}

The Centers for Disease Control and Prevention's (CDC) *Learn the Signs. Act Early.* program, funded the American Academy of Pediatrics (AAP) to convene an expert working group to revise its developmental surveillance checklists. The goals of the group were to identify evidence-informed milestones to include in CDC checklists, clarify when most children can be expected to reach a milestone (to discourage a wait-and-see approach), and support clinical judgment regarding screening between recommended ages. Subject matter experts identified by the AAP established 11 criteria for CDC milestone checklists, including using milestones most children ($\geq 75\%$) would be expected to achieve by specific health supervision visit ages and those that are easily observed in natural settings. A database of normative data for individual milestones, common screening and evaluation tools, and published clinical opinion was created to inform revisions. Application of the criteria established by the AAP working group and adding milestones for the 15- and 30-month health supervision visits resulted in a 26.4% reduction and 40.9% replacement of previous CDC milestones. One third of the retained milestones were transferred to different ages; 67.7% of those transferred were moved to older ages. Approximately 80% of the final milestones had normative data from ≥ 1 sources. Social-emotional and cognitive milestones had the least normative data. These criteria and revised checklists can be used to support developmental surveillance, clinical judgment regarding additional developmental screening, and research in developmental surveillance processes. Gaps in developmental data were identified particularly for social-emotional and cognitive milestones.

The American Academy of Pediatrics (AAP) recommends developmental surveillance and screening to identify children with developmental delays or disabilities (DDs) early, help to ensure timely interventions, and improve outcomes.¹ Developmental surveillance is a longitudinal process that involves eliciting concerns, taking a developmental history based on milestone attainment, observing milestones and other behaviors, examining the

child, and applying clinical judgment during health supervision visits (HSVs). Developmental screening involves the use of validated screening tools at specific ages or when surveillance reveals a concern.¹ Diagnostic evaluations are conducted, typically by developmental specialists, to further evaluate and diagnose DDs in children deemed at risk through surveillance and screening processes.

abstract



^aNational Center on Birth Defects and Developmental Disabilities, Centers for Disease Control and Prevention, Atlanta, Georgia; ^bEagle Global Scientific, LLC, San Antonio, Texas; ^cDivision of Developmental-Behavioral Pediatrics, Medical University of South Carolina, Charleston, South Carolina; ^dDivision of Developmental Pediatrics, Department of Pediatrics, The University of Tennessee Health Science Center, Memphis, Tennessee; ^eVermont Child Health Improvement Program, Department of Pediatrics, Larner College of Medicine, The University of Vermont, Burlington, Vermont; ^fCenter on Human Development (Professor Emerita), University of Oregon, Eugene, Oregon; ^gMetroHealth Medical Center, Case Western Reserve University, Cleveland, Ohio; ^hAmerican Academy of Pediatrics, Itasca, Illinois; ⁱKennedy Krieger Institute, Baltimore, Maryland; ^jJohns Hopkins School of Medicine, Baltimore, Maryland; and ^kNational Center on Birth Defects and Developmental Disabilities (retired), Centers for Disease Control and Prevention, Atlanta, Georgia * Contributed equally as co-senior authors.

Drs Zubler and Wiggins conceptualized and designed the study, collected data and performed the initial analyses, drafted the initial manuscript, and reviewed and revised the manuscript; Drs Lipkin, Macias, Whitaker, Squires, Pajek and Shaw critically reviewed and revised the manuscript for important intellectual content; Ms Wolf, Ms Slaughter, Ms Broughton, Ms Gerndt, and Ms Mlodoch planned and performed the research and reviewed and revised the manuscript; and all authors approved the final manuscript as submitted and agree to be accountable for all aspects of the work.

The contents are those of the author(s) and do not necessarily represent the official views of, nor an endorsement by American Academy of Pediatrics, Centers for Disease Control and Prevention/US Department of Health and Human Services, or the US government.

DOI: <https://doi.org/10.1542/peds.2021-052138>

Accepted for publication Dec 9, 2021

Address correspondence to Jennifer M. Zubler, MD, National Center on Birth Defects and Developmental Disabilities, Centers for Disease Control and Prevention, 4770 Buford Hwy NE, MS S106-4, Atlanta, GA 30341. E-mail: wyy4@cdc.gov

To cite: Zubler JM, Wiggins LD, Macias MM, et al. Evidence-Informed Milestones for Developmental Surveillance Tools. *Pediatrics*. 2022;149(3):e2021052138

Developmental surveillance is family centered to promote conversations and trusting relationships wherein families can express concerns.² Surveillance involves clinical judgment about when a child may be at risk for delays and when additional developmental screening might be warranted. Milestone lists help to guide developmental surveillance, but those used for surveillance, unlike screening and evaluation tools, are not validated. Typically, lists of milestones are uncited, are based on clinical opinion, and/or report the average or median age a milestone should be achieved. Moreover, ages specified for individual milestones are inconsistent across sources.^{3,4}

Lists that cite average or median ages at which children achieve milestones provide insight into typical development but do not provide clarity for parents, pediatricians, and other early childhood professionals (ECPs) about when to be concerned or when additional screening might be helpful.⁴ For example, lists based on median (50th percentile) age milestones might encourage a wait-and-see⁵ approach because half of children are not expected to achieve the milestone by that age. In CDC focus groups, parents of children with disabilities reported delays in identification because they were told to wait, that children develop differently, and that some take longer than others. Milestone lists need to support developmental surveillance and clinical judgment on when additional developmental screening could better assess risk for developmental delays.

In 2004, the CDC's *Learn the Signs. Act Early.* program developed free developmental surveillance milestone checklists that included developmental warning signs for parents, pediatricians, and ECPs; messaging to "act early" by

addressing concerns; and developmental tips/activities. These materials were developed to help parents to recognize typical development, elicit parents' concerns about their child's development, improve discussions between parents and professionals about a child's development, and support universal developmental screening at recommended ages and additional screenings when there are concerns. The milestones were adapted from *Caring for Your Baby and Young Child: Birth to Age 5* (5th ed) to align with recommended HSVs.⁶ Like most milestone lists, the original sources of the milestones were uncited, and adaptations were based mainly on clinical opinion, not on empirically informed evidence.

Based on 15 years of use, 3 areas for improving the checklists were identified by the CDC. First, criteria for checklists used for surveillance needed to be established to evaluate the existing CDC checklists. Second, milestones within checklists would represent milestones above the 50th percentile^{3,4,7,8} to ensure that most children would achieve the milestone by a given age. Lastly, new checklists for the 15- and 30-month HSVs were needed to complete the series 2 months to 5 years of age to improve integration of developmental surveillance across early childhood HSVs. This article presents the results of these revisions.

METHODS

The AAP Systems of Services for Children and Youth With Special Health Care Needs team identified and convened 8 subject matter experts (SMEs) in different fields of child development. The group included developmental-behavioral, neurodevelopmental, and general pediatricians; child and developmental psychologists; and a professor of special education and

early intervention. All SMEs had graduate training and experience in research methodology and medical decision-making and clinical experience in developmental surveillance, screening, and evaluation. One SME was an editor of and 2 contributed to the AAP's *Bright Futures: Guidelines for Health Supervision of Infants, Children, and Adolescents* (4th ed).⁹ Two SMEs were lead authors of the AAP's 2020 clinical report, "Promoting Optimal Development: Identifying Infants and Young Children With Developmental Disorders Through Developmental Surveillance and Screening"¹; another had long-time experience as a developer of screening tools.

Criteria Identification

The SMEs developed 11 criteria for CDC surveillance milestones and tools (Table 1). SMEs nominated the criteria on the basis of their clinical experience and use of CDC surveillance materials. Nominations were discussed, and those that were unanimously agreed upon were included as criteria. Of note, the SMEs agreed that milestones should be easily observed in natural settings and $\geq 75\%$ of children would be expected to achieve a milestone at a given age. The strategy ($\geq 75\%$) was chosen to support clinical judgment regarding performing additional developmental screenings, with validated screening tools as a next step to assess a child's risk for developmental delays. This strategy may also prevent a wait-and-see approach because most children of the same age would be expected to achieve the milestone. To reduce confusion about when to be concerned, developmental warning signs were eliminated because not achieving milestones that most children ($\geq 75\%$) are expected to achieve similarly warrants more in-

TABLE 1 Criteria for Developmental Milestones and Surveillance Tools.

1. Milestones are included at the age most ($\geq 75\%$) children would be expected to demonstrate the milestone
2. Eliminate "warning signs"^a
3. Are easy for families of different social, cultural, and ethnic backgrounds to observe and use
4. Are able to be answered with yes, not yet, or not sure
5. Use plain language, avoiding vague terms like may, can, and begins
6. Are organized in developmental domains
7. Show progression of skills with age, when possible
8. Milestones are not repeated across checklists
9. Include open-ended questions
10. Include information for developmental promotion
11. Include information on how to act early if there are concerns

Criteria developed by SMEs.

^a Milestones listed separately within CDC materials with parent messaging to act early if child has not attained them.

depth surveillance and consideration for developmental screening.

Milestone Identification

Milestones for possible inclusion in CDC surveillance materials were identified by the SMEs. Existing CDC milestones were automatically included for evaluation and were the foundation for the revised checklists. Other milestones were identified by a literature review and commonly used developmental resources.

Literature Review

A broad literature search was conducted in March 2019 using MEDLINE, PsychInfo, and ERIC databases. Search terms were developed in collaboration with the CDC librarian after discussion of the objective of the literature review and comprised the following: (1) milestone, normative (data, table, range, value) or age (range, appropriate) AND (2) child development or infant development AND (3) percentile, psychometrics, predictive values, red flags, warning signs, assessment, monitor, delay, or reference (standard, values).

An article was included for evaluation if it was written in English, contained evidence that supported at least one normed individual developmental milestone or included published clinical opinion (ie, consensus milestones) that children exhibit the milestone

by a specific age, and limited to children aged ≤ 5 years. Articles were excluded if evidence was limited to special populations (eg, preterm infants) or risk factors. Non-peer-reviewed articles, dissertations, and books were also excluded.

Articles were evaluated by the first 2 authors to determine if they met inclusion criteria. Articles chosen by both authors, either on initial review or after additional discussion and agreement between these 2 authors, had milestone data extracted for review by the SME group.

Developmental Resources

SMEs nominated several additional resources for evaluation, including parent resources, professional teaching resources, and commonly used screening and diagnostic evaluation tools (Table 2). Inclusion criteria for these resources were availability in English and contained evidence that supported at least one individual developmental milestone or included published clinical opinion (ie, consensus milestones) that children exhibit the milestone by a specific age. All nominations were accepted for additional review.

CDC Milestones

Existing CDC milestones served as a foundation for identifying milestones and were automatically evaluated.

Milestone Evaluation

The following data were extracted from existing CDC milestones, articles selected for review, and developmental resources: individual milestones, supporting references, and any normative data or published clinical opinion for the milestone. Milestones with conflicting data (eg, age at which most children should achieve the milestone) were flagged for additional discussion.

4A comprehensive database was created with information extracted from CDC milestones and resources for individual milestones from ages 2 months to 5 years, aligned with AAP HSV ages, and shared with the SME group. Original data sources, such as articles that met inclusion criteria and nominated developmental resources, were also shared with the SME group. SMEs collaborated through an in-person meeting, 6 virtual meetings, and e-mail reviews of decision summaries from January to September 2019.

During meetings, the SMEs discussed categorizing milestones into 4 developmental domains: (1) social emotional, (2) language/communication, (3) cognitive, and (4) motor. These domains were previously used in CDC materials and could help parents to learn about different areas of child development (eg, social-emotional

TABLE 2 Additional Developmental Resources Reviewed

Parent Resources	Educational/Training Resources	Developmental Screening Tools	Diagnostic Evaluation Tools ^a
AAP Bright Futures Previsit Questionnaires ¹⁰	AAP Bright Futures guidelines (4th ed) ⁹ AAP <i>Pediatrics in Review</i> articles ^{15–21}	Ages & Stages Questionnaires (3rd ed) ²²	Bayley Scales of Infant and Toddler Development (3rd ed) ²⁷
American Speech-Language-Hearing Association development charts ¹¹		Ages & Stages Questionnaires: Social-Emotional ²³	Beery-Buktenica Developmental Test of Visual-Motor Integration (6th ed) ²⁸
AAP brochure “Is Your One-Year-Old Communicating With You?” ¹²		Modified Checklist for Autism in Toddlers, Revised ²⁴	BRIGANCE Early Childhood Screens III ²⁹
CDC <i>Learn the Signs. Act Early.</i> checklists ¹³		Parents’ Evaluation of Developmental Status With Developmental Milestones ²⁵	The Capute Scales: Cognitive Adaptive Test/Clinical Linguistic and Auditory Milestone Scale ³⁰
FIRST WORDS Project 16 × 16 ¹⁴		Survey of Well-Being in Young Children ²⁶	MacArthur-Bates Communicative Development Inventories (2nd ed) ³¹
			Mullen Scales of Early Learning ³²
			Peabody Developmental Motor Scales (2nd ed) ³³
			Preschool Language Scale-5 ³⁴

^a Diagnostic evaluation tools were cross referenced when there was lack of agreement supporting a milestone or age of a milestone across other data sources. Not all milestones were cross referenced with diagnostic resources.

skills in addition to language/communication skills). Because milestones often represent skills across several domains, they were placed in the domain in which the SME group believed that parents would most likely identify them. For example, reciprocal play skills involving other people were categorized as social-emotional, and other play skills were categorized as cognitive.

Milestones were reviewed for 2 to 3 ages at a time (eg, 2, 4, and 6 months) so that skill progression could be considered. Milestones were reviewed by the SME group on the basis of the criteria outlined in Table 1. First, SMEs considered milestones with normative data that supported achievement by $\geq 75\%$ of children at a particular age. Next, they considered milestones from screening and diagnostic tools. Finally, SMEs considered milestones that were based on published clinical opinion. Each of these data sources was cross referenced with the others to get a sense of the evidence base available to support inclusion of a milestone in revised CDC checklists.

If there was disagreement across sources or lack of evidence supporting a milestone for a specific age, additional research was conducted, and additional evidence was then evaluated by the SMEs according to the developed criteria. This included reviewing diagnostic evaluation tools (Table 2) and/or conducting a separate PubMed search for a specific milestone or a related skill. For example, additional evidence was sought for age-specific development of gestures like lifting arms to be picked up, waving bye-bye, and blowing a kiss.

SMEs used the available evidence base and their clinical experience to determine if and at what age a milestone would be included in the revised checklists. Only milestones with unanimous agreement were included.

During the evaluation process, SMEs denoted milestones they included for surveillance but believed that additional research could improve age placement, quality, and/or quantity of supporting evidence or better capture the underlying developmental construct of the milestone. Finally, the SMEs

simplified the milestones and added examples to try to improve understanding. The CDC team then reviewed milestones from a cultural and health communication perspective (eg, family friendly, fifth- to sixth-grade reading level), and SMEs reviewed those changes again to ensure that the milestone still represented the developmental construct being assessed.

With the use of back-translation methods, the milestones were translated into Spanish by CDC Multilingual Services and reviewed by 2 native Spanish-speaking pediatricians. Cognitive testing with a diverse sample of parents located in different regions of the United States provided feedback on reliability and clarity, which led the SMEs to make additional changes to wording. Results of cognitive testing will be published separately.

RESULTS

Of the 1027 articles generated from the literature review, 34 met inclusion criteria. Of those, 24 contained normative data, and 10 contained published clinical opinion for ≥ 1 milestones. Six additional articles were found through

independent searches to evaluate individual milestones when SMEs determined that the milestone was appropriate but insufficient or conflicting evidence supported its use at a specific HSV age. Of the articles with normative data, 14 (58.3%) described populations within non-English-speaking countries.

Tables 3 to 6 list the milestones that SMEs included in CDC checklists after critical evaluation and unanimous decision. They are presented by domain with HSV age, references that support the inclusion of the milestone at that age, and whether the milestone was an existing or new CDC milestone. Supporting references are divided into normative data, developmental screening and evaluation tools, and published clinical opinions.

Previously, CDC had 216 milestones across 10 checklists. With the addition of 15- and 30-month checklists and the evidence review process, 159 milestones were included across 12 checklists. This represented a reduction of 57 (26.4%) CDC milestones, with the average number of milestones per checklist decreasing from 22 to 13.

Of the final 159 milestones that met the evaluation criteria, 94 (59.1%) were based on CDC original milestones and 65 (40.9%) were added on the basis of the milestone identification and evaluation process. One third of the 94 retained CDC milestones were moved to a different age on the basis of the criterion that $\geq 75\%$ of children would be expected to achieve the milestone by that age. When moved, 21 of those 31 milestones were transferred to an older age. More than half (56.5%) of the original 216 milestones were eliminated (Supplemental Table 7) on the basis of SME criteria, including 25 that were duplicated across checklists at

different ages. For example, “Tries to use things the right way, like a phone, cup, or book” was on both the 12- and 18-month checklists; this milestone was placed only at 15 months on the basis of supporting evidence. Additionally, eliminating vague terms, such as may or begins, resulted in moving and changing milestones; for example, “Begins to pass things from one hand to another” was removed from the list of milestones at age 6 months and was included as “Moves things from one hand to her other hand” at age 9 months.

The 1-, 2-, and 3-year-old checklists had the greatest decrease in the number of milestones, with a $\geq 50\%$ reduction. Approximately half of the aggregate loss for these ages was due to moving milestones to the new 15- and 30-month checklists. When combining all new and revised checklists for 1 to 3 years, the number of milestones was reduced by 25.7%, consistent with the 26.4% reduction of milestones across the other age ranges.

All 4 domains had a reduction in number of milestones. Cognitive milestones decreased by 34 (50.7%), social emotional decreased by 16 (27.5%), language decreased by 4 (9.1%), and motor decreased by 3 (6.4%). Social-emotional and cognitive domain milestones were the least likely to have normative data available. The social-emotional domain had 25 (59.5%) milestones with 0 to 1 normed references and 8 (19.0%) with ≥ 3 normed references. The cognitive domain had 19 (57.6%) milestones with 0 to 1 normed references and 8 (24.2%) with ≥ 3 references. In contrast, language and motor domains had 11 (27.5%) and 10 (22.7%) milestones with 0 to 1 normed references and 21 (52.5%) and 30 (68.1%) milestones, respectively, with ≥ 3 normed references.

Of the final 159 milestones, 127 (79.9%) were retained or added on the basis of normative data, whereas 32 (20.1%) milestones were included on the basis of screening and evaluation tools, published clinical opinion, and SME opinion. Of milestones with normative data, 32 had normative data from only 1 resource. The SMEs believed that 22 (13.8%) of the final milestones were candidates for additional research, as noted in Tables 3–6; this subset includes the milestone “Sings, dances, or acts for you,” the only milestone included or retained without supporting evidence on the basis of SME opinion.

Of the 77 developmental warning signs listed on the old CDC checklists, 59 (76.6%) had a corresponding milestone on the new checklists.

DISCUSSION

The CDC checklists support developmental surveillance and other important components of the early identification process by pediatricians and other ECPs, including developmental promotion, parent education and engagement, communication of developmental progress and concerns, and developmental screening.^{731,5,9,73–76} However, variability across surveillance resources, including the CDC’s, can create confusion regarding what constitutes a concern and when developmental screening between recommended ages might be warranted.^{3,4,38,77}

An expert working group convened by AAP sought to improve CDC surveillance tools by enhancing conversations among pediatricians, ECPs, and families regarding childhood development and guiding clinical judgment on when to conduct developmental screening between recommended ages. These tools are not intended to replace

TABLE 3 Social-Emotional Milestones With Supporting Normative Data, Evaluation Tools, and Published Clinical Opinion References

Social-Emotional Milestones	Age	CDC or New	Normative Data ^a	Source	
				Developmental Screening and Evaluation Tools ^b	Published Clinical Opinion ^c
Calms down when spoken to or picked up ^d	2 mo	New	Ertem et al (35)	—	ASHA (11), Bright Futures (36), Dosman et al (8), Sharp et al (37)
Looks at your face	2 mo	CDC	Ertem et al (35), Sheldrick and Perrin (38)	—	Bright Futures (36), Dosman et al (8), Scharf et al (18)
Seems happy to see you when you walk up to her	2 mo	New	Ertem et al (35), Sheldrick and Perrin (38), Thalagala (39)	—	—
Smiles when you talk to or smile at her	2 mo	CDC	Bhawe et al (40), Ertem et al (35), Lejarraga et al (41), Thalagala (39)	ASQ-3 (22), PEDS-DM (25)	ASHA (11)
Smiles on his own to get your attention	4 mo	CDC	Ertem et al (35)	ASQ-3, PEDS-DM	Bright Futures (36)
Chuckles (not yet a full laugh) when you try to make her laugh	4 mo	New	Accardo and Capute (30), Bhawe et al (40), Ertem et al (35), Sheldrick and Perrin (38)	ASQ-3, PEDS-DM	ASHA (11), Bellman et al (42), Bright Futures (36)
Looks at you, moves, or makes sounds to get or keep your attention	4 mo	New	Ertem et al (35)	PEDS-DM	—
Knows familiar people	6 mo	CDC	Bhawe et al (40), Ertem et al (35)	ASQ-3	Bright Futures (36), Dosman et al (8), Gerber et al (20), Scharf et al (18)
Likes to look at himself in the mirror	6 mo	CDC	Bhawe et al (40), Lejarraga et al (41)	ASQ-3	Bright Futures (36)
Laughs	6 mo	New	Accardo and Capute (30), Bhawe et al (40), Ertem et al (35), Sheldrick and Perrin (38)	ASQ-3, PEDS-DM	Bellman et al (42), Bright Futures (36), Dosman et al (8)
Is shy, clingy, or fearful around strangers	9 mo	CDC	Ertem et al (43), Kumar et al (44), Lancaster et al (45)	—	Gerber et al (20), Scharf et al (18)
Shows several facial expressions, like happy, sad, angry, and surprised ^e	9 mo	New	Thalagala (39)	—	Gerber et al (20), Scharf et al (18)
Looks when you call his name	9 mo	CDC	Gladstone et al (46), Sheldrick and Perrin (38)	ASQ-3, PEDS-DM	Bright Futures (36), Dosman et al (8),

TABLE 3 Continued

Social-Emotional Milestones	Age	CDC or New	Normative Data ^a	Source	
				Developmental Screening and Evaluation Tools ^b	Published Clinical Opinion ^c
Reacts when you leave (looks, reaches for you, or cries) ^d	9 mo	CDC	Ertem et al (43)	—	Gerber et al (20), Scharf et al (18), Dosman et al (8), Gerber et al (20), Scharf et al (18), Bellman et al (42), Bright Futures (36), Dosman et al (8), Gerber et al (20), Scharf et al (18)
Smiles or laughs when you play peek-a-boo	9 mo	CDC	Ertem et al (35), Lejamaga et al (41), Sheldrick and Perrin (38), Thalagala (39)	PEDS-DM	Bellman et al (42), Bright Futures (36), Dosman et al (8), Gerber et al (20), Scharf et al (18)
Plays games with you, like pat-a-cake	12 mo	CDC	Ertem et al (35), Fenson et al (47), Sheldrick and Perrin (38)	—	Bellman et al (42), Bright Futures (36), Gerber et al (20), Scharf et al (18)
Copies other children while playing, like taking toys out of a container when another child does ^d	15 mo	CDC	—	—	Bright Futures (36), Dosman et al (8)
Shows you an object that he likes	15 mo	New	Crais et al (48)	ASQ-3	Dosman et al (8), First Words (14), Gerber et al (20), Scharf et al (18)
Claps when excited	15 mo	New	Crais et al (48), Kwon et al (49)	—	—
Hugs stuffed doll or other toy	15 mo	New	Crais et al (48), Fenson et al (47)	—	Colson and Dworkin (16), Johnson and Blasco (15)
Shows you affection (hugs, cuddles, or kisses you)	15 mo	CDC	Ertem et al (43)	ASQ-SE-2 (23), Bayley III (27)	Johnson and Blasco (15), Vaughan (17)
Moves away from you, but looks to make sure you are close by	18 mo	CDC	—	ASQ-SE-2	Dosman et al (8), Gerber et al (20), Scharf et al (18)
Points to show you something interesting	18 mo	CDC	—	Bayley III, MCHAT-R (24), PEDS-DM	Bellman et al (42), Bright Futures (36), Dosman et al (8), Gerber et al (18)

TABLE 3 Continued

Social-Emotional Milestones	Age	CDC or New	Normative Data ^a	Source	
				Developmental Screening and Evaluation Tools ^b	Published Clinical Opinion ^c
Puts hands out for you to wash them	18 mo	New	Gladstone et al (46)	—	(20), Scharf et al (18)
Looks at a few pages in a book with you ^d	18 mo	New	Ertem et al (35)	ASQ-SE-2	Behrman et al (50), Colson and Dworkin (16), Vaughan (17)
Helps you dress him by pushing arm through sleeve or lifting up foot	18 mo	CDC	—	ASQ-3, PEDS-DM	Bright Futures (36), Dosman et al (8), Johnson and Blasco (15)
Notifies when others are hurt or upset, like pausing or looking sad when someone is crying	24 mo	New	—	PEDS-DM	Dosman et al (8), Scharf et al (18)
Looks at your face to see how to react in a new situation	24 mo	New	—	ASQ-SE-2, MCHAT-R, PEDS-DM	Bright Futures (36), Dosman et al (8), Johnson and Blasco (15)
Plays next to other children and sometimes plays with them	30 mo	CDC	Lancaster et al (45), Lansdown et al (51)	ASQ-SE-2, PEDS-DM	Bright Futures (36), Dosman et al (8), Gerber et al (20), Johnson and Blasco (15), Scharf et al (18)
Shows you what she can do by saying, "Look at me!"	30 mo	New	Sheldrick and Perrin (38)	—	—
Follows simple routines when told, like helping to pick up toys when you say, "It's clean-up time."	30 mo	New	—	ASQ-SE, PEDS-DM	Colson and Dworkin (16), Gerber et al (20), Johnson and Blasco (15), Scharf et al (18), Vaughan (17)
Calms down within 10 min after you leave her, like at child care drop off ^e	3 y	CDC	—	—	Dosman et al (8), Johnson and Blasco (15)
Notifies other children and joins them to play	3 y	New	Lansdown et al (51)	ASQ-SE-2, PEDS-DM	Dosman et al (8), Gerber et al (20), Johnson and Blasco (15), Scharf et al (18)
	4 y	CDC	Ertem et al (43)	ASQ-3, PEDS-DM	

TABLE 3 Continued

Social-Emotional Milestones	Age	CDC or New	Source		Published Clinical Opinion ^c
			Normative Data ^a	Developmental Screening and Evaluation Tools ^b	
Pretends to be something else during play (teacher, superhero, dog)	4 y	CDC	Gladstone et al (46)	—	Behrman et al (50), Bright Futures (36), Dosman et al (8)
Asks to go play with children if none are around, like “Can I play with Alex?” ^d	4 y	CDC	—	ASQ-3, PEDS-DM	Dosman et al (8)
Comforts others who are hurt or sad, like hugging a crying friend ^d	4 y	New	—	ASQ-SE-2	—
Avoids danger, like not jumping from tall heights at the playground	4 y	New	Gladstone et al (46), Haitiwanger and Coستر (52)	PEDS-DM	Gerber et al (20), Scharf et al (18)
Likes to be a “helper”	4 y	New	Gladstone et al (46)	ASQ-SE-2, PEDS-DM	—
Changes behavior on the basis of where she is (place of worship, library, playground)	4 y	New	Gladstone et al (46), Haitiwanger and Coستر (52), Kumar et al (44), Sheldrick and Perrin (38)	ASQ-3, PEDS-DM	Bright Futures (36), Colson and Dworkin (16), Dosman et al (8), Gerber et al (20), Scharf et al (18)
Follows rules or takes turns when playing games with other children	5 y	CDC	Gladstone et al (46), Haitiwanger and Coستر (52)	—	—
Sings, dances, or acts for you ^d	5 y	CDC	—	PEDS-DM	—
Does simple chores at home, like matching socks or clearing the table after eating	5 y	New	Gladstone et al (46), Haitiwanger and Coستر (52)	—	—

ASHA, American Speech-Language-Hearing Association; ASQ-3, Ages & Stages Questionnaires (3rd ed); ASQ-SE-2, Ages & Stages Questionnaires: Social-Emotional (2nd ed); Bayley III, Bayley Scales of Infant and Toddler Development (3rd ed); MCHAT-R, Modified Checklist for Autism in Toddlers, Revised; PEDS-DM, Parents’ Evaluation of Developmental Status With Developmental Milestones. —, Empty cells indicate that type/category of data was not available to support inclusion of that milestone.

^a Provides baseline distribution data for age of attainment of a milestone for a given population.

^b Some commonly used developmental screening and diagnostic evaluation tools (not all available tools are represented).

^c Uncited on the basis of clinical opinion and/or report of the average or median age that a milestone should be achieved.

^d SME agreement that additional research would be beneficial.

validated screening tools but instead to promote optimal child development and encourage professionals to act early through surveillance and screening as outlined in the AAP's clinical report.¹ Results indicate substantial changes were made developing and applying criteria for surveillance milestones and tools, adding 15- and 30-month checklists, and incorporating evidence-informed milestones.

Clinicians have used attainment of developmental milestones for almost a century⁷⁸ to determine if a child is developing typically. Pediatricians have reported increasing use of milestone checklists from 53.0% in 2002 to 89.6% in 2016, in addition to their report of increased developmental screening.⁷⁹ However, these surveillance milestone checklists are likely based on published clinical opinion due to the lack of published normative milestone data and the lack of citations of original sources on checklists. The domain tables generated from this work linking developmental milestones to empirically informed evidence and published clinical opinion could improve training of professionals and methods for surveillance,^{4,7,8} such as incorporation of the CDC's open access milestones/checklists into electronic health records.

Milestone checklists used in surveillance are intended to prompt conversations, review developmental history and progress, and elicit concerns. The CDC checklists should not replace universal developmental screening, provide a risk categorization, or diagnose DDs. To determine if concerns about milestones should prompt a more in-depth developmental history, observation, and examination along with consideration for assessing actual risk by screening, the SMEs

recommended that most children ($\geq 75\%$) should be expected to achieve milestones by a given age.

SMEs believed that using 50th percentile milestones for surveillance would not support clinical decision-making for developmental screening because only half of children would be expected to achieve an individual milestone by a given age. Using milestones that 85% or 90% of children would be expected to achieve may limit opportunities for additional screening for too many children at risk for developmental delays. A $\geq 75\%$ criterion was thus agreed upon to balance informed clinical decision-making regarding developmental screening and provide opportunities to identify children at risk for delays as soon as possible. Pediatricians are encouraged to follow AAP recommendations to screen a child, using validated screening tools, when surveillance reveals a concern or anytime a concern is raised, to identify and refine the risk for developmental delays.¹

The criterion (Table 1) for milestones representing those that most children ($\geq 75\%$) would be expected to achieve eliminated the need for the CDC's previous warning signs because most are now represented as milestones. Using this strategy, SMEs agreed that a child not meeting a milestone should be considered for screening similar to children demonstrating warning signs. Less than a quarter of warning signs could not be replaced with an evidence-informed milestone. Examples not represented as new milestones include subjective items such as "shows extreme behavior (unusually fearful, aggressive, shy or sad)" and items that may be better recognized during a physical examination, such as "stiff or tight muscles." Additional features of the surveillance tools

may identify those types of concerns. For example, the act early message was retained, and new open-ended questions were added to encourage parents to ask about other concerns.

Use of this same strategy is intended to discourage the wait-and-see approach and could prevent worry for children older than the average age of attainment of a milestone but not likely to be at risk for delays.^{1,5} It could also eliminate the confusion families had with milestones and warning signs on the same checklist and repetition of milestones across checklists.

The application of additional criteria (Table 1) to improve clarity in the revised milestones may help parents, pediatricians, and other ECPs to recognize when missing milestones might indicate the need for developmental screening. Although milestones typically span several developmental domains, SMEs believed that keeping domain labels could improve awareness of the less commonly known social-emotional and cognitive development milestones of young children. Showing progression of a skill, like walking, could demonstrate how earlier milestones lay the foundation for later ones. Adding the open-ended question, "Is there anything your child is doing or is not doing that concerns you?" may encourage parents to discuss the quality with which a child exhibits a milestone, milestones not listed within the sample, and atypical behaviors difficult to capture through a list.⁴² Finally, including information on developmental promotion and acting early may empower families to support their child's development and to ask about screening if they have concerns.

Surveillance of social-emotional and cognitive milestones supports

TABLE 4 Language and Communication Milestones With Supporting Normative Data, Evaluation Tools, and Published Clinical Opinion References

Language/Communication Milestones	Age	CDC or New	Normative Data ^a	Source	
				Developmental Screening and Evaluation Tools ^b	Published Clinical Opinion ^c
Makes sounds other than crying	2 mo	CDC	Dosman et al (8), ^d Sheldrick and Perrin (38)	ASQ-3 (22), PEDS-DM (25)	Blackwell and Baker (53), Bright Futures (36), Gerber et al (20), Scharf et al (18)
Reacts to loud sounds ^e	2 mo	New	—	—	Accardo and Capute (30), ASHA (11), Bellman et al (42), Bright Futures (36)
Makes sounds like “oooo” and “aahh” (cooling)	4 mo	CDC	Accardo and Capute (30), Ertem et al (35)	ASQ-3	ASHA (11), Bright Futures (36), Dosman et al (8), Gerber et al (20), Scharf et al (18)
Makes sounds back when you talk to him	4 mo	New	Ertem et al (35), Den Ouden et al (54), Kumar et al (44)	ASQ-3	Bellman et al (42), Dosman et al (8), Gerber et al (20), Johnson and Blasco (15), Scharf et al (18)
Turns head toward the sound of your voice	4 mo	CDC	Accardo and Capute (30), Bhawe et al (40), Den Ouden et al (54), Sheldrick and Perrin (38)	—	Bellman et al (42), Dosman et al (8), Gerber et al (20), Scharf et al (18)
Takes turns making sounds with you	6 mo	CDC	Kumar et al (44), Lejarraga et al (41)	—	Dosman et al (8), Gerber et al (20), Scharf et al (18)
Blows “raspberries” (sticks tongue out and blows)	6 mo	New	Accardo and Capute (30)	—	Blackwell and Baker (53), Gerber et al (20), Scharf et al (18)
Makes squealing noises	6 mo	New	—	ASQ-3	Bellman et al (42), Blackwell and Baker (53), Gerber et al (20), Scharf et al (18)
Makes different sounds like “mamamama” and “babababa”	9 mo	CDC	Accardo and Capute (30), Bhawe et al (40), Den Ouden et al (54), Lancaster et al (45), Lejarraga et al (41), Sheldrick and Perrin (38)	ASQ-3, PEDS-DM	ASHA (11), Bright Futures (36)
Lifts arms up to be picked up	9 mo	New	Ertem et al (35), Fenson et al (47), Gladstone et al (46), Kwon et al (49), Sheldrick and Perrin (38)	—	ASHA (11), Bright Futures (36), Dosman et al (8), First Words (14), Gerber et al (20), Scharf et al (18)

TABLE 4 Continued

Language/Communication Milestones	Age	CDC or New	Normative Data ^a	Source	
				Developmental Screening and Evaluation Tools ^b	Published Clinical Opinion ^c
Waves "bye-bye"	12 mo	CDC	Accardo and Capute (30), Crais et al (48), Den Ouden et al (54), Ertem et al (35), Fenson et al (47), Kwon et al (49)	—	ASHA (11), Bright Futures (36), First Words (14), Gerber et al (20), Scharf et al (18)
Calls a parent "mama" or "dada" or another special name	12 mo	CDC	Accardo and Capute (30), Crais et al (48), Sheldrick and Perrin (38)	—	AAP (12), Bright Futures (36), Gerber et al (20), Scharf et al (18)
Understands "no" (pauses briefly or stops when you say it)	12 mo	CDC	Accardo and Capute (30), Ertem et al (35), Gladstone et al (46), Lancaster et al (45), Lejarraga et al (41), Thalagala (39)	Bayley III (27)	Blackwell and Baker (53), Dosman et al (8), Gerber et al (20), Scharf et al (18)
Tries to say 1 or 2 words besides mama or dada, like "ba" for ball or "da" for dog	15 mo	New	Gladstone et al (46), Lancaster et al (45), Tamis-LeMonda et al (55)	ASQ-3, PEDS DM	Bright Futures (36), Johnson and Blasco (15), Sharp and Hillenbrand (37)
Looks at a familiar object when you name it	15 mo	CDC	Ertem et al (35), Lancaster et al (45), Sheldrick and Perrin (38)	ASQ-3	Gerber et al (20), Scharf et al (18)
Follows directions given with both a gesture and words. For example, he gives you a toy when you hold out your hand and say, "Give me the toy."	15 mo	New	Accardo and Capute (30), Ertem et al (35), Gladstone et al (46)	ASQ-3	Bright Futures (36), Johnson and Blasco (15)
Points to ask for something or to get help	15 mo	CDC	Ertem et al (43), Kwon et al (49), Lancaster et al (45)	ASQ-3	AAP (12), Gerber et al (20), Johnson and Blasco (15), Scharf et al (18)
Tries to say ≥3 words besides mama or dada	18 mo	CDC	Accardo and Capute (30), Ertem et al (35), Gladstone et al (46), Lancaster et al (45)	ASQ-3	Bellman et al (42), Blackwell and Baker (53), Bright Futures (36), Dosman et al (8), Gerber et al (20), Scharf et al (18)

TABLE 4 Continued

Language/ Communication Milestones	Age	CDC or New	Normative Data ^a	Source	
				Developmental Screening and Evaluation Tools ^b	Published Clinical Opinion ^c
Follows 1-step directions without any gestures, like giving you the toy when you say, "Give it to me." Points to things in a book when you ask, for example, "Where is the bear?"	18 mo	CDC	(45), Sheldrick and Perrin (38) Accardo and Capute (30), Ertem et al (35), Gladstone et al (46), Lancaster et al (45), Sheldrick and Perrin (38)	—	ASHA (11), Dosman et al (8), Gerber et al (20), Scharf et al (18)
Says at least 2 words together, like "More milk."	24 mo	CDC	Accardo and Capute (30), Den Ouden (54), Ertem et al (45), Gladstone et al (46), Sheldrick and Perrin (38)	ASQ-3	ASHA (11), Bellman et al (42), Blackwell and Baker (53), Bright Futures (36), Gerber et al (20), Johnson and Blasco (15), Scharf et al (18) AAP (12), ASHA (11), Bellman et al (42), Bright Futures (36), Dosman et al (8), Gerber et al (20), Scharf et al (18)
Points to at least 2 body parts when you ask him to show you Uses more gestures than just waving and pointing, like blowing a kiss or nodding yes Says ~50 words	24 mo	CDC	Accardo and Capute (30), Muluk et al (56), Sheldrick and Perrin (38) Fenson et al (47), Kwon et al (49)	ASQ-3, PEDS-DM	ASHA (11), Bellman et al (42), Blackwell and Baker (53), Bright Futures (36), Johnson and Blasco (15), Dosman et al (8), First Words (13)
Says ≥2 words, with 1 action word, like "Doggie run."	30 mo	New	Accardo and Capute (30), Lancaster et al (45), Tarnis-Lemonda et al (55)	ASQ-3	AAP (11), Bright Futures (36), Gerber et al (20), Johnson and Blasco (15), Scharf et al (18)
Names things in a book when you point and ask, "What is this?"	30 mo	New	Accardo and Capute (30), Ertem et al (35), Lancaster et al (45), Tarnis-Lemonda et al (55) Lancaster et al (45), Sheldrick and Perrin (38)	ASQ-3, Bayley III	Blackwell and Baker (53), Dosman et al (8), Gerber et al (20), Johnson and Blasco (15), Scharf et al (18) Gerber et al (20), Scharf et al (18)

TABLE 4 Continued

Language/ Communication Milestones	Age	CDC or New	Normative Data ^a	Source	
				Developmental Screening and Evaluation Tools ^b	Published Clinical Opinion ^c
Says words like I, me, or we	30 mo	CDC	Accardo and Capute (30), Ertem et al (35), Lancaster et al (45), Sheldrick and Perrin (38)	ASQ-3, Bayley III	Bright Futures (36), Colson and Dworkin (16), Gerber et al (20), Johnson and Blasco (15), Scharf et al (18)
Talks with you in conversation using at least 2 back- and-forth exchanges ^e	3 y	CDC	—	Bayley III	—
Asks who, what, where, or why questions, like “Where is mommy/ daddy?”	3 y	New	Ertem et al (43), Nair and Russell (57), Sheldrick and Perrin (38)	—	ASHA (11), Colson and Dworkin (16)
Says what action is happening in a picture when asked, like running, eating, or playing	3 y	New	Gladstone et al (46)	ASQ-3, Bayley III	Gerber et al (20), Knobloch et al (58), Scharf et al (18)
Says first name when asked	3 y	CDC	Gladstone et al (46), Haltiwanger and Coster (52), Lansdown et al (51), Lejarraga et al (41), Sheldrick and Perrin (38)	ASQ-3	Gerber et al (20), Knobloch et al (58), Scharf et al (18)
Talks well enough for others to understand, most of the time	3 y	CDC	Ertem et al (35), Gladstone et al (46), Lancaster et al (45)	Bayley III	ASHA (11), Bellman et al (42), Blackwell and Baker (53), Bright Futures (36), Dosman et al (8), Gerber et al (20), Johnson and Blasco (15), Scharf et al (18)
Says sentences with ≥ 4 words ^e	4 y	New	—	ASQ-3	ASHA (11)
Says some words from a song, story, or nursery rhyme ^e	4 y	CDC	—	—	Colson and Dworkin (16)
Talks about at least 1 thing that happened during	4 y	New	—	—	ASHA (11)

TABLE 4 Continued

Language/ Communication Milestones	Age	ODC or New	Normative Data ^a	Source	
				Developmental Screening and Evaluation Tools ^b	Published Clinical Opinion ^c
his day, like "I played soccer." ^e Answers simple questions, like "What is a coat for," or "What is a crayon for?"	4 y	New	Ertem et al (43), Gladstone et al (46), Lejarraaga et al (41), Nair and Russell (57), Sheldrick and Perrin (38)	ASQ-3	ASHA (11), Gerber et al (20), Scharf et al (18)
Tells a story she heard or made up with at least 2 events, like a cat stuck in a tree and a firefighter saving it	5 y	ODC	McCabe and Rollins (59), Sheldrick and Perrin (38)	—	Gerber et al (20), Scharf et al (18)
Answers simple questions about a book or story after you read or tell it to him ^d	5 y	New	—	—	ASHA (11)
Keeps a conversation going with >3 back-and-forth exchange ^e	5 y	New	—	—	ASHA (11)
Uses or recognizes simple rhymes (bat-cat, ball-tail)	5 y	New	—	PEDS-DM	ASHA (11), Gerber et al (20), Scharf et al (18)

ASHA, American Speech-Language-Hearing Association; ASQ-3, Ages & Stages Questionnaires (3rd ed); Bayley III, Bayley Scales of Infant and Toddler Development (3rd ed); PEDS-DM, Parents' Evaluation of Developmental Status With Developmental Milestones. —, Empty cells indicate that type/category of data was not available to support inclusion of that milestone.

^a Provides baseline distribution data for age of attainment of a milestone for a given population.

^b Some commonly used developmental screening and diagnostic evaluation tools (not all available tools are represented).

^c Uncited on the basis of clinical opinion and/or report the average or median age that a milestone should be achieved.

^d Dosman et al⁸ contained both published clinical opinion and normative data.

^e SME agreement that additional research would be beneficial.

early identification of children with intellectual and social-emotional disorders, including autism spectrum disorder. In our review, milestones included in the social-emotional domain had the fewest normed references per milestone, followed by the cognitive domain, language domain, and motor domain. When social-emotional milestones were available, they were often self-help rather than social engagement and emotional regulation skills. These results highlight the need for additional research on social-emotional milestones. For example, can social-emotional function be assessed using milestones during developmental surveillance at HSVs, and what factors influence the timing and development of social-emotional skills? Despite relatively more evidence in the language domain, SMEs believed that more research is needed in the intersection of the social-emotional and language domains to recognize and monitor the development of social language in young children.

Limitations

There were limitations with the revision process. SMEs' opinions determined which HSV was most appropriate for a milestone, given existing evidence and the goal that most children ($\geq 75\%$) would be expected to achieve it by that age. For example, if there was evidence that 50% of children reach a milestone at 11 months and 90% reach it at 16 months, it was placed on the 15-month checklist. Eighty percent of the revised milestones had normative data to support their inclusion on a specific HSV age checklist. However, it was necessary to use data such as published clinical opinion to represent milestones in all domains across all ages.

Regardless of supporting data type, all revised milestones had unanimous SME support for their inclusion in a surveillance tool. When normative data existed, limitations remained in the availability of milestones that met other criteria. For example, sometimes it was difficult to find naturally observable milestones compared with milestones demonstrated during standardized testing. Although the milestones are evidence-informed using international sources and cognitive testing with parents was done, there is no certainty that these new milestones will resonate with all families or that they are the most relevant milestones for developmental surveillance. These checklists and others have not been tested to see whether they indeed lead to appropriate developmental screening between recommended ages and improve early identification.

There are other gaps in developmental surveillance research. Best practices for conducting the recommended 6 components¹ of surveillance are not available. There are limitations in the use of milestones, even evidence-informed ones, as the sole component of developmental surveillance. Milestones/milestone checklists can support some components, such as taking a developmental history and eliciting concerns, but surveillance also involves observation, examination, education, communication, and clinical decision-making. Moreover, trusting relationships may develop through the longitudinal process of surveillance and improve parents' confidence in sharing concerns and pediatricians'

confidence to assess concerns raised.

CONCLUSIONS

Early identification and intervention for the 1 in 6 children with DDs have been shown to improve outcomes.⁸⁰⁻⁸⁴ However, less than a quarter of children with DDs receive early intervention services before age 3 years,^{85,86} and most children with emotional, behavioral, and developmental conditions, other than autism spectrum disorder, do not receive services before age 5 years.⁸⁷ Developmental surveillance is an important part of early identification^{1,88} and facilitates education, communication, and relationship building among parents, pediatricians, and ECPs. Research in early identification has focused primarily on developmental screening, not surveillance. Improvements in surveillance tools and processes could help to identify concerns and support clinical judgment regarding developmental screening to allow more timely referral to early intervention services and additional evaluation. The methods described herein led to substantial revisions of CDC resources to better support developmental surveillance. Best practices for surveillance and improvements in surveillance tools could be supported by additional research on individual normed milestones, particularly social-emotional and cognitive milestones; how parents, pediatricians, and ECPs learn and understand milestones; which milestones are most likely to lead to appropriate screening; whether the use of milestones that most children ($\geq 75\%$) would be expected to achieve is appropriate for surveillance; whether categorizing milestones into

TABLE 5 Cognitive Milestones With Supporting Normative Data, Evaluation Tools, and Published Clinical Opinion References

Cognitive Milestones	Age	CDC or New	Normative Data ^a	Source	
				Developmental Screening and Evaluation Tools ^b	Published Clinical Opinion ^c
Watches you as you move	2 mo	CDC	Accardo and Capute (30), Atkinson et al (60), Bhawe et al (40), Ertem et al (35), Sheldrick and Perrin (38), Kumar et al (44)	ASQ-3 (22)	—
Looks at a toy for several seconds	2 mo	New		—	Accardo and Capute (30), Bellman et al (42)
If hungry, opens mouth when she sees breast or bottle	4 mo	New	Camruth and Skinner (61), Thalagala (39)	ASQ-3, PEDS-DM (25)	Gerber et al (20), Johnson and Blasco (15), Scharf et al (18)
Looks at his hands with interest	4 mo	New	Den Ouden et al (54), Ertem et al (35), Lejarraaga et al (41)	—	Bellman et al (41), Gerber et al (20), Johnson and Blasco (15), Scharf et al (18)
Puts things in her mouth to explore them	6 mo	CDC	Ertem et al (35)	ASQ-3	Gerber et al (20), Scharf et al (18)
Reaches to grab a toy he wants	6 mo	CDC	Den Ouden et al (54), Kumar et al (44)	ASQ-3	Dosman et al (8), Gerber et al (20), Johnson and Blasco (15), Scharf et al (18)
Closes lips to show she does not want more food ^d	6 mo	New	—	PEDS-DM	—
Looks for objects when dropped out of sight (like his spoon or toy)	9 mo	New	Accardo and Capute (30), Ertem et al (35)	—	Bright Futures (36), Dosman et al (8), Gerber et al (20), Johnson and Blasco (15), Scharf et al (18)
Bangs 2 things together	9 mo	CDC	Ertem et al (35), Gladstone et al (46), Sheldrick and Perrin (38)	ASQ-3	Bellman et al (42), Bright Futures (36), Gerber et al (20), Scharf et al (18)
	12 mo	CDC	Accardo and Capute (30), Gladstone et al	ASQ-3	

TABLE 5 Continued

Cognitive Milestones	Age	CDC or New	Normative Data ^a	Source	
				Developmental Screening and Evaluation Tools ^b	Published Clinical Opinion ^c
Puts something in a container, like a block in a cup	12 mo	CDC	(46), Lancaster et al (45), Thalagala (39)	—	Bright Futures (36), Gerber et al (20), Scharf et al (18)
Looks for things he sees you hide, like a toy under a blanket	15 mo	CDC	Accardo and Capute (30), Ertem et al (35), Gladstone et al (46)	—	Bright Futures (36), Gerber et al (20), Scharf et al (18)
Tries to use things the right way, like a phone, cup, or book ^d	15 mo	New	—	Beery VMI (28)	Dosman et al (8), Johnson and Blasco (15)
Stacks at least 2 small objects, like blocks	18 mo	CDC	Den Ouden et al (54), Fenson et al (47)	ASQ-3, PEDS-DM	Bellman et al (42), Dosman et al (8), Gerber et al (20), Scharf et al (18)
Copies you doing chores, like sweeping with a broom	18 mo	CDC	Ertem et al (43), Gladstone et al (46), Lancaster et al (45)	ASQ-3	Bellman et al (42), Dosman et al (8), Gerber et al (20), Johnson and Blasco (15), Scharf et al (18)
Plays with toys in a simple way, like pushing a toy car	24 mo	New	Kimmerle et al (62)	—	—
Holds something in 1 hand while using the other hand, for example, holding a container and taking the lid off ^d	24 mo	CDC	Ertem et al (35)	ASQ-3	Bright Futures (36), Colson and Dworkin (16), Dosman et al (8), Gerber et al (20), Scharf et al (18)
Tries to use switches, knobs, or buttons on a toy	24 mo	New	—	PEDS-DM	—
Plays with >1 toy at the same time, like putting toy food on a toy plate	30 mo	New	—	ASQ-3	—

TABLE 5 Continued

Cognitive Milestones	Age	CDC or New	Normative Data ^a	Source		Published Clinical Opinion ^c
				Developmental Screening and Evaluation Tools ^b		
Uses things to pretend, like feeding a block to a doll as if it were food	30 mo	New	Lancaster et al (45)	ASQ-3	—	Bright Futures (36), Dosman et al (8), Gerber et al (20), Scharf et al (18)
Shows simple problem-solving skills, like standing on a small stool to reach something	30 mo	CDC	Accardo and Capute (30), Ertem et al (35), Gladstone et al (46) Lancaster et al (45)	Bayley III (27)	—	AAP (12), Bright Futures (36), Dosman et al (8), Gerber et al (20), Scharf et al (18), Knobloch et al (58)
Follows 2-step instructions, for example, "Put the toy down and close the door."	30 mo	New	Accardo and Capute (30), Sheldrick and Perrin (38)	—	—	—
Shows that he knows at least 1 color, like pointing to a red crayon when you ask, "Which one is red?"	3 y	CDC	Accardo and Capute (30), Egan and Brown (63)	ASQ-3	—	Behrman et al (50), Bellman et al (42), Bright Futures (36), Colson and Dworkin (16), Dosman et al (8), Gerber et al (20), Scharf et al (18)
Draws a circle when you show him how	3 y	New	Haltiwanter and Coster (52)	ASQ-SE (23)	—	—
Avoids touching hot objects, like a stove, when you warn her ^d	4 y	CDC	Accardo and Capute (30), Lejarraaga et al (41), Sheldrick and Perrin (38)	ASQ-3, Bayley III, PEDS-DM	—	Gerber et al (20), Scharf et al (18)
Names a few colors of items	4 y	CDC	—	—	—	Gerber et al (20), Scharf et al (18)
Tells what comes next in a well-known story ^d	4 y	CDC	Lejarraaga et al (41),	ASQ-3, PEDS-DM	—	—

TABLE 5 Continued

Cognitive Milestones	Age	CDC or New	Normative Data ^a	Source	
				Developmental Screening and Evaluation Tools ^b	Published Clinical Opinion ^c
Draws a person with ≥ 3 body parts					
Counts to 10	5 y	CDC	—	ASQ-3	Bright Futures (36), Gerber et al (20), Scharf et al (18), Dosman et al (8), Gerber et al (20), Scharf et al (18)
Names some numbers between 1 and 5 when you point to them	5 y	CDC	Williams and Lerner (64)	PEDS-DM	ASHA (11), Gerber et al (20), Scharf et al (18)
Uses words about time, like yesterday, tomorrow, morning, or night	5 y	CDC	Sheldrick and Perrin (38)	PEDS-DM	Colson and Dworkin (16), Gerber et al (20), Scharf et al (18)
Pays attention for 5–10 min during activities, for example, during story time or making arts and crafts (screen time does not count) ^d	5 y	New	—	ASQ-SE	—
Writes some letters in her name	5 y	CDC	Sheldrick and Perrin (38)	ASQ-3, PEDS-DM	Dosman et al (8), Gerber et al (20), Scharf et al (18)
Names some letters when you point to them	5 y	New	Williams and Lerner (64)	ASQ-3	Gerber et al (20), Scharf et al (18)

ASHA, American Speech-Language-Hearing Association; ASQ-3, Ages & Stages Questionnaires (3rd ed); ASQ-SE, Ages & Stages Questionnaires: Social-Emotional; Bayley III, Bayley Scales of Infant and Toddler Development (3rd ed); Beery VMI, Beery-Buktenica Developmental Test of Visual-Motor Integration; PEDS-DM, Parents' Evaluation of Developmental Status With Developmental Milestones. —, Empty cells indicate that type/category of data was not available to support inclusion of that milestone.

^a Provides baseline distribution data for age of attainment of a milestone for a given population.

^b Some commonly used developmental screening and diagnostic evaluation tools (not all available tools are represented).

^c Uncited on the basis of clinical opinion and/or report the average or median age that a milestone should be achieved.

^d SME agreement that additional research would be beneficial.

TABLE 6 Motor Milestones With Supporting Normative Data, Evaluation Tools, and Published Clinical Opinion References

Motor Milestones	Age	CDC or New	Normative Data ^a	Source	
				Developmental Screening and Evaluation Tools ^b	Published Clinical Opinion ^c
Holds head up when on tummy	2 mo	CDC	Accardo and Capute (30), Carruth and Skinner (61), Den Ouden et al (54)	—	Bright Futures (36), Dosman et al (8), Gerber et al (20), Scharf et al (18), Bright Futures (36)
Moves both arms and both legs	2 mo	New	—	ASQ-3 (22)	Bright Futures (36)
Opens hands briefly	2 mo	New	Accardo and Capute (30), Ertem et al (35), Lejarraga et al (41)	ASQ-3	Bright Futures (36)
Holds head steady without support when you are holding her	4 mo	CDC	Ertem et al (35), Lejarraga et al (41), Sheldrick and Perrin (38)	PEDS-DM (25)	Bright Futures (36), Gerber et al (20), Scharf et al (18)
Holds a toy when you put it in his hand	4 mo	CDC	Dosman et al (8) ^d	—	Bellman et al (42), Gerber et al (20), Scharf et al (18)
Uses her arm to swing at toys	4 mo	CDC	Bhawe et al (40), Dosman et al (8), Ertem et al (43), Kumar et al (44)	—	Gerber et al (20), Scharf et al (18), Scharf et al (18)
Brings hands to mouth	4 mo	CDC	Den Ouden et al (54), Ertem et al (35), Lejarraga et al (41), Sheldrick and Perrin (38)	—	Bright Futures (36), Dosman et al (8)
Pushes up onto elbows/forearms when on tummy	4 mo	CDC	Accardo and Capute (30), Lejarraga et al (41), Thalagala (39)	—	Gerber et al (20), Scharf et al (18)
Rolls from tummy to back	6 mo	CDC	Accardo and Capute (30), Den Ouden et al (54), Dosman et al (8), Ertem et al (35)	ASQ-3	Gerber et al (20), Scharf et al (18)
Pushes up with straight arms when on tummy	6 mo	CDC	Accardo and Capute (30), Carruth and Skinner (61), Thalagala (39)	—	Gerber et al (20), Scharf et al (18)
Leans on hands to support himself when sitting	6 mo	CDC	Accardo and Capute (30), Carruth and Skinner (61),	ASQ-3	Gerber et al (20), Scharf et al (18)

TABLE 6 Continued

Motor Milestones	Age	CDC or New	Normative Data ^a	Source	
				Developmental Screening and Evaluation Tools ^b	Published Clinical Opinion ^c
Gets to a sitting position by herself	9 mo	CDC	Dosman et al (8), Ertem et al (35), Kitsao-Wekulo et al (65), Lejarraaga et al (41) Accardo and Capute (30), Bhawe et al (40), Gejewska et al (66), Sheldrick and Perrin (38)	—	Gerber et al (20), Scharf et al (18)
Sits without support	9 mo	CDC	Accardo and Capute (30), Bhawe et al (40), Cox et al (67), Dosman et al (8), Ertem et al (35), Gladstone et al (46), Lancaster et al (44), Lejarraaga et al (41)	—	Bellman et al (42), Gerber et al (20), Scharf et al (18)
Uses fingers to “rake” food toward himself	9 mo	New	Accardo and Capute (30), Carruth and Skinner (61), Ertem et al (43), Gladstone et al (46), Sheldrick and Perrin (38)	ASQ-3	Bright Futures (36), Gerber et al (20), Scharf et al (18)
Moves things from 1 hand to her other hand	9 mo	CDC	Accardo and Capute (30), Carruth and Skinner (61), Cox et al (67), Den Ouden et al (54), Gladstone et al (46), Kitsao-Wekulo et al (65), Lancaster et al (44), Lejarraaga et al (41), Sheldrick and Perrin (38), Thalagala (39)	ASQ-3	Bright Futures (36), Dosman et al (8), Gerber et al (20), Scharf et al (18)
Pulls up to stand	12 mo	CDC	Accardo and Capute (30), Bhawe et al (40), Den Ouden et al (54), Dosman et al (8), Ertem et al	—	Bellman et al (42), Gerber et al (20), Scharf et al (18)

TABLE 6 Continued

Motor Milestones	Age	CDC or New	Normative Data ^a	Source	
				Developmental Screening and Evaluation Tools ^b	Published Clinical Opinion ^c
Walks, holding onto furniture	12 mo	CDC	(43), Gladstone et al (46), Lancaster et al (45), Lejarraaga et al (41), World Health Organization (68) Accardo and Capute (30), Ertem et al (43), Lejarraaga et al (41), World Health Organization (68) Gladstone et al (46), Lancaster et al (45)	—	Bellman et al (42), Gerber et al (20), Scharf et al (18)
Drinks from a cup without a lid, as you hold it	12 mo	New	Gladstone et al (46), Lancaster et al (45)	ASQ-3, PEDS-DM	Bright Futures (36), Gerber et al (20), Johnson and Blasco (15), Scharf et al (18)
Picks things up between thumb and pointer finger, like small bits of food	12 mo	CDC	Ertem et al (43), Gladstone et al (46), Kumar et al (44), Lancaster et al (45), Lejarraaga et al (41)	—	Bright Futures (36), Dosman et al (8), Gerber et al (20), Scharf et al (18)
Takes a few steps on his own	15 mo	CDC	Ertem et al (43), Gladstone et al (46), Noller and Ingrisano (69)	—	Bright Futures (36)
Uses fingers to feed herself some food	15 mo	New	Camruth and Skinner (61), Dosman et al (8), Kumar et al (44), Lejarraaga et al (41)	—	Bright Futures (36), Gerber et al (20), Scharf et al (18)
Walks without holding onto anyone or anything	18 mo	CDC	Accardo and Capute (30), Bhawe et al (40), Gladstone et al (46), Kitsao-Wekulo et al (65), Lancaster et al (45), Lejarraaga et al (41), Sheldrick and Perrin (38)	ASQ-3, PEDS-DM	Bellman et al (42), Bright Futures (36), Dosman et al (8)
Scribbles	18 mo	CDC	Accardo and Capute (30), Ertem et al (43), Kitsao-Wekulo et al (65), Lancaster et al (45), Lejarraaga et al (41), Sheldrick and Perrin (38)	ASQ-3	Bright Futures (36), Dosman et al (8), Gerber et al (20), Johnson and

TABLE 6 Continued

Motor Milestones	Age	CDC or New	Normative Data ^a	Source		Published Clinical Opinion ^c
				Developmental Screening and Evaluation Tools ^b		
Drinks from a cup without a lid and may spill sometimes	18 mo	CDC	et al (45), Lansdown et al (51), Lejarraaga et al (41) Den Ouden et al (53), Ertem et al (43), Gladstone et al (46), Lancaster et al (45)	ASQ-3		Blasco (15), Scharf et al (18) Bright Futures (36), Dosman et al (8), Gerber et al (20), Johnson and Blasco (15), Scharf et al (18) Bright Futures (36), Gerber et al (20), Scharf et al (18)
Feeds herself with her fingers	18 mo	New	Carruth and Skinner (61), Dosman et al (8), Kumar et al (44), Lejarraaga et al (41)	—		Bellman et al (42), Bright Futures (70), Dosman et al (8), Gerber et al (20), Johnson and Blasco (15), Scharf et al (18)
Tries to use a spoon	18 mo	New	Ertem et al (43), Gladstone et al (46)	ASQ-3		Bellman et al (42), Bright Futures (70), Dosman et al (8), Gerber et al (20), Johnson and Blasco (15), Scharf et al (18)
Climbs on and off a couch or chair without help	18 mo	CDC	Carruth and Skinner (61), Kitsao-Wekulo et al (65), Lancaster et al (45), Lejarraaga et al (41)	—		Bright Futures (36), Colson and Dworkin (16), Gerber et al (20), Johnson and Blasco (15), Scharf et al (18)
Kicks a ball	24 mo	CDC	Ertem et al (43), Gladstone et al (46), Lancaster et al (45), Lejarraaga et al (41), Sheldrick and Perrin (38), Thalagala (39)	ASQ-3		Bellman et al (42), Bright Futures (36), Colson and Dworkin (16), Dosman et al (8), Gerber et al (20), Johnson and Blasco (15), Scharf et al (18)
Runs	24 mo	CDC	Accardo and Capute (30), Gladstone et al (46), Sheldrick and Perrin (38)	ASQ-3		Bellman et al (42), Bright Futures (36), Colson and Dworkin (16), Dosman et al (8),

TABLE 6 Continued

Motor Milestones	Age	CDC or New	Normative Data ^a	Source		Published Clinical Opinion ^c
				Developmental Screening and Evaluation Tools ^b		
Walks (not climbs) up a few stairs with or without help	24 mo	CDC	Bhave et al (40), Ertem et al (43), Sheldrick and Perrin (38)	ASQ-3, PEDS-DM		Gerber et al (20), Scharf et al (18), Bellman et al (42), Bright Futures (70), Colson and Dworkin (16), Dosman et al (8), Gerber et al (20), Johnson and Blasco (15), Scharf et al (18)
Eats with a spoon	24 mo	CDC	Dosman et al (8), Ertem et al (43), Gladstone et al (46), Muluk et al (56)	—		Bright Futures (36), Colson and Dworkin (16), Dosman et al (8), Gerber et al (20), Johnson and Blasco (15), Scharf et al (18)
Uses hands to twist things, like turning doorknobs or unscrewing lids	30 mo	CDC	Lancaster et al (45)	MSEL (32)		Bright Futures (36), Johnson and Blasco (15)
Takes some clothes off by himself, like loose pants or an open jacket	30 mo	CDC	Dosman et al (8), Ertem et al (43), Gladstone et al (46)	ASQ-3, PEDS-DM		Bellman et al (42), Bright Futures (36), Gerber et al (20), Johnson and Blasco (15), Scharf et al (18)
Jumps off the ground with both feet	30 mo	New	Gladstone et al (46), Kitsao-Wekulo et al (65), Lejarraga et al (41), Sheldrick and Perrin (38)	ASQ-3		Bright Futures (36), Colson and Dworkin (16), Gerber et al (20), Johnson and Blasco (15), Scharf et al (18)
Turns book pages, 1 at a time, when you read to her	30 mo	CDC	—	MSEL		Bright Futures (36), Knobloch et al (58)
Strings items together, like large beads or macaroni	3 y	New	Gladstone et al (46)	ASQ-3		Gerber et al (20), Knobloch et al

TABLE 6 Continued

Motor Milestones	Age	CDC or New	Normative Data ^a	Source	
				Developmental Screening and Evaluation Tools ^b	Published Clinical Opinion ^c
Puts on some clothes by himself, like loose pants or a jacket	3 y	CDC	Dosman et al (8), Ertem et al (35)	ASQ-3, PEDS-DM	(58), Scharf et al (18), Bellman et al (42), Bright Futures (36), Gerber et al (20), Johnson and Blasco (15), Scharf et al (18)
Uses a fork	3 y	CDC	Dosman et al (8)	ASQ-3	Bellman et al (42), Bright Futures (36)
Catches a large ball most of the time	4 y	CDC	—	ASQ-3	Bright Futures (36), Gerber et al (20), Scharf et al (18)
Serves himself food or pours water, with adult supervision	4 y	CDC	—	ASQ-3	Gerber et al (20), Scharf et al (18)
Unbuttons some buttons	4 y	New	Ohtoshi et al (71)	—	Bright Futures (36), Gerber et al (20), Scharf et al (18)
Holds crayon or pencil between fingers and thumb (not in a fist)	4 y	New	Egan and Brown (63), Ertem et al (43)	ASQ-3	Bright Futures (36), Dosman et al (8), Knobloch et al (58)
Buttons some buttons	5 y	New	Ohtoshi et al (71)	ASQ-3	Bright Futures (36), Gerber et al (20), Scharf et al (18)
Hops on 1 foot	5 y	CDC	Gladstone et al (46), Kitsao-Wekulo et al (65), Nair and Russell (57)	ASQ-3	Bellman et al (42), Bright Futures (36), Gerber et al (20), Scharf et al (18)

ASQ-3, Ages & Stages Questionnaires (3rd ed); MSEL, Mullen Scales of Early Learning; PEDS-DM, Parents' Evaluation of Developmental Status With Developmental Milestones. —, Empty cells indicate that type/category of data was not available to support inclusion of that milestone.

^aProvides baseline distribution data for age of attainment of a milestone for a given population.

^bSome commonly used developmental screening and diagnostic evaluation tools (not all available tools are represented).

^cUncited, based on clinical opinion, and/or report the average or median age that a milestone should be achieved.

^dDosman et al⁸ contained both published clinical opinion and normative data.

developmental domains is helpful; and whether cultural differences exist in surveillance milestones and processes.^{1,3–5,7} Nevertheless, based on review of milestone data and clinical experience, the SMEs agreed that most typically developing children would achieve the developmental constructs represented. To our knowledge, this attempt is the first to align empirically informed milestones on parent-completed surveillance tools with objectively defined

criteria agreed upon by SMEs. The CDC milestones and checklists can be used in continued efforts to improve developmental surveillance.

ACKNOWLEDGMENTS

We thank Mary Cogswell, RN, DrPH, for her review of earlier versions of the manuscript and pediatricians Natalia Benza, MD, and Jose O. Rodriquez, MD, MBA, for their review of the Spanish milestone translations.

ABBREVIATIONS

AAP: American Academy of Pediatrics
 CDC: Centers for Disease Control and Prevention
 DD: developmental delay or disability
 ECP: early childhood professional
 HSV: health supervision visit
 SME: subject matter expert

Copyright © 2022 by the American Academy of Pediatrics

FUNDING: This project is supported by the Centers for Disease Control and Prevention of the US Department of Health and Human Services as part of a financial assistance award totaling \$100,000, with 100% funded by the Centers for Disease Control and Prevention/US Department of Health and Human Services.

CONFLICT OF INTEREST DISCLOSURES: Dr Squires is a developer of the *Ages & Stages Questionnaires* and receives royalties from Brookes Publishing, the company that publishes this tool; the other authors have indicated they have no conflicts of interest relevant to this article to disclose.

REFERENCES

- Lipkin PH, Macias MM; Council on Children With Disabilities, Section on Developmental and Behavioral Pediatrics. Promoting optimal development: identifying infants and young children with developmental disorders through developmental surveillance and screening. *Pediatrics*. 2020;145(1):e20193449
- Foy JM, Green CM, Earls MF; Committee on Psychosocial Aspects of Child and Family Health, Mental Health Leadership Work Group. Mental health competencies for pediatric practice. *Pediatrics*. 2019;144(5):e20192757
- Wilkinson CL, Wilkinson MJ, Lucarelli J, et al. Quantitative evaluation of content and age concordance across developmental milestone checklists. *J Dev Behav Pediatr*. 2019;40(7):511–518
- Sices L. Use of developmental milestones in pediatric residency training and practice: time to rethink the meaning of the mean. *J Dev Behav Pediatr*. 2007;28(1):47–52
- Raspa M, Levis DM, Kish-Doto J, et al. Examining parents' experiences and information needs regarding early identification of developmental delays: qualitative research to inform a public health campaign. *J Dev Behav Pediatr*. 2015;36(8):575–585
- Shelov, SP, Altmann, T, *Caring for Your Baby and Young Child: Birth to Age 5*, New and Revised 5th ed, American Academy of Pediatrics, Itasca, IL, 2009
- Sheldrick RC, Schlichting LE, Berger B, et al. Establishing new norms for developmental milestones. *Pediatrics*. 2019;144(6):e20190374
- Dosman CF, Andrews D, Goulden KJ. Evidence-based milestone ages as a framework for developmental surveillance. *Paediatr Child Health*. 2012;17(10):561–568
- Hagan JFSJ, Duncan PM, eds. *Bright Futures: Guidelines for Health Supervision of Infants, Children, and Adolescents*, 4th ed. Itasca, IL: American Academy of Pediatrics; 2017
- Shaw J, Hagan JJ, Shepard M, et al. *Bright Futures Tool and Resource Kit*. Itasca, IL: American Academy of Pediatrics; 2019
- American Speech-Language-Hearing Association. How Does Your Child Hear and Talk? Available at: <https://www.asha.org/public/speech/development/chart>. Accessed August 26, 2021
- American Academy of Pediatrics. Is Your One-Year-Old Communicating With You? [brochure]. Itasca, IL: American Academy of Pediatrics; 2005
- National Center on Birth Defects and Developmental Disabilities, Centers for Disease Control and Prevention. Tools for Tracking Milestones. Available at: <https://www.cdc.gov/ncbddd/actearly/freematerials.html>. Accessed August 26, 2021
- FIRST WORDS Project. 16 by 16. Available at: <https://firstwordsproject.com/about-16by16>. Accessed August 26, 2021
- Johnson CP, Blasco PA. Infant growth and development. *Pediatr Rev*. 1997;18(7):224–242
- Colson ER, Dworkin PH. Toddler development. *Pediatr Rev*. 1997;18(8):255–259
- Vaughan VC III. Assessment of growth and development during infancy and early childhood. *Pediatr Rev*. 1992;13(3):88–97
- Scharf RJ, Scharf GJ, Stroustrup A. Developmental milestones. *Pediatr Rev*. 2016;37(1):25–37, quiz 38, 47
- Gerber RJ, Wilks T, Erdie-Lalena C. Developmental milestones 3: social-emotional development. *Pediatr Rev*. 2011;32(12):533–536
- Gerber RJ, Wilks T, Erdie-Lalena C. Developmental milestones: motor

- development. *Pediatr Rev*. 2010;31(7):267–276, quiz 277
21. Wilks T, Gerber RJ, Erdie-Lalena C. Developmental milestones: cognitive development. *Pediatr Rev*. 2010;31(9):364–367
 22. Squires J, Bricker D. *Ages & Stages Questionnaires*. Baltimore, MD: Brookes Publishing; 2009
 23. Squires J, Bricker D, Twombly E. *Ages & Stages Questionnaires: Social-Emotional*. Baltimore, MD: Brookes Publishing; 2002
 24. Robins M, Fein D, Barton M. *The Modified Checklist for Autism in Toddlers, Revised With Follow-Up (M-CHAT-R/F)*. 2009
 25. Glascoe F, Robertshaw N. *PEDS: Developmental Milestones*. Nashville, TN: Ellsworth & Vandermeer Press; 2018
 26. Tufts Medical Center. The Survey of Children Well-being of Young Children. Available at: <https://www.tuftschildrenshospital.org/the-survey-of-wellbeing-of-young-children/overview>. Accessed August 26, 2021
 27. Bayley N. *The Bayley Scales of Infant and Toddler Development*. 3rd ed. Minneapolis, MN: Pearson; 2006
 28. Beery KE, Buktenica NA, Beery NA. *The Beery–Buktenica Developmental Test of Visual–Motor Integration: Administration, Scoring, and Teaching Manual*. Minneapolis, MN: Pearson; 2010
 29. French B. *BRIGANCE Early Childhood Screens III*. North Billerica, MA: Curriculum Associates; 2013
 30. Accardo P, Capute A. *The Capute Scales: Cognitive Adaptive Test/Clinical Linguistic and Auditory Milestone Scale*. Baltimore, MD: Brookes Publishing; 2005
 31. Fenson LMV, Thal D, Dale P, Reznick JS, Bates E. *MacArthur-Bates Communicative Development Inventories Baltimore*. Baltimore, MD: Brookes Publishing; 2007
 32. Mullen E. *Mullen Scales of Early Learning*. Bloomington, MN: Pearson; 1995
 33. Folio MR, Fewell RR. *Peabody Developmental Motor Scales*. Austin, TX: PRO-ED; 2000
 34. Zimmerman IL, Steiner VG, Pond RA. *The Preschool Language Scale-5*. San Antonio, TX: Pearson; 2011
 35. Ertem IO, Krishnamurthy V, Mulaudzi MC, et al. Similarities and differences in child development from birth to age 3 years by sex and across four countries: a cross-sectional, observational study. *Lancet Glob Health*. 2018;6(3):e279–e291
 36. Lipkin P, Macias M. *Developmental Milestones for Developmental Surveillance at Preventive Care Visits. Bright Futures: Guidelines for Health Supervision of Infants, Children, and Adolescents*. 4th ed. Itasca, IL: American Academy of Pediatrics; 2017
 37. Sharp HM, Hillenbrand K. Speech and language development and disorders in children. *Pediatr Clin North Am*. 2008;55(5):1159–1173, viii
 38. Sheldrick RC, Perrin EC. Evidence-based milestones for surveillance of cognitive, language, and motor development. *Acad Pediatr*. 2013;13(6):577–586
 39. Thalagala N. Windows of achievement for development milestones of Sri Lankan infants and toddlers: estimation through statistical modelling. *Child Care Health Dev*. 2015;41(6):1030–1039
 40. Bhave A, Bhargava R, Kumar R. Development and validation of a new Lucknow Development Screen for Indian children aged 6 months to 2 years. *J Child Neurol*. 2010;25(1):57–60
 41. Lejarraaga H, Pascucci MC, Krupitzky S, et al. Psychomotor development in Argentinean children aged 0-5 years. *Paediatr Perinat Epidemiol*. 2002;16(1):47–60
 42. Bellman M, Byrne O, Sege R. Developmental assessment of children. *BMJ*. 2013;346(7891):e8687
 43. Ertem IO, Krishnamurthy V, Mulaudzi MC, et al. Validation of the International Guide for Monitoring Child Development demonstrates good sensitivity and specificity in four diverse countries. *Acta Paediatr*. 2019;108(6):1074–1086
 44. Kumar R, Iyengar SD, Bhasin S, Gupta I, Kumar V. A normative study of child development using a culture-appropriate test battery in rural Haryana, India. *J Trop Pediatr*. 1995;41(1):38–42
 45. Lancaster GA, McCray G, Kariger P, et al. Creation of the WHO Indicators of Infant and Young Child Development (IYCD): metadata synthesis across 10 countries. *BMJ Glob Health*. 2018;3(5):e000747
 46. Gladstone M, Lancaster GA, Umar E, et al. The Malawi Developmental Assessment Tool (MDAT): the creation, validation, and reliability of a tool to assess child development in rural African settings. *PLoS Med*. 2010;7(5):e1000273
 47. Fenson L, Dale PS, Reznick JS, Bates E, Thal DJ, Pethick SJ. Variability in early communicative development. *Monogr Soc Res Child Dev*. 1994;59(5):1–173, discussion 174–185
 48. Crais ER, Watson LR, Baranek GT. Use of gesture development in profiling children's prelinguistic communication skills. *Am J Speech Lang Pathol*. 2009;18(1):95–108
 49. Kwon AY, Vallotton CD, Kiegelmann M, Wilhelm KH. Cultural diversification of communicative gestures through early childhood: a comparison of children in English-, German-, and Chinese-speaking families. *Infant Behav Dev*. 2018;50:328–339
 50. Behrman R, Kiegelman R, Arvin A, et al. *Nelson's Textbook of Pediatrics*. Philadelphia, PA: WB Saunders; 1996
 51. Lansdown RG, Goldstein H, Shah PM, et al. Culturally appropriate measures for monitoring child development at family and community level: a WHO collaborative study. *Bull World Health Organ*. 1996;74(3):283–290
 52. Haltiwaner J, Coster W. *A Normative Study of Development in Context: Growth toward Independence in Social Function Skills of Young Children*. Washington, DC: National Institute on Disability and Rehabilitation Research ED/OSERS; 1991:29
 53. Blackwell PB, Baker BM. Estimating communication competence of infants and toddlers. *J Pediatr Health Care*. 2002;16(1):29–35
 54. Den Ouden L, Rijken M, Brand R, Verloove-Vanhorick SP, Ruys JH. Is it correct to correct? Developmental milestones in 555 “normal” preterm infants compared with term infants. *J Pediatr*. 1991;118(3):399–404
 55. Tamis-Lemonda CS, Bornstein MH, Kahana-Kalman R, Baumwell L, Cyphers L. Predicting variation in the timing of language milestones in the second year: an events history approach. *J Child Lang*. 1998;25(3):675–700
 56. Muluk NB, Bayoğlu B, Konuşkan B, Anlar B. Milestones of language development in Turkish children. *B-ENT*. 2013;9(4):299–306
 57. Nair MKC, Russell PS. Development and normative validation of Developmental Assessment Tool for Anganwadis for 3- to 4-year-old children (DATA-II). *J Clin Epidemiol*. 2013;66(1):23–29
 58. Knobloch H, Stevens F, Malone A. *Manual of Developmental Diagnosis: The Administration*

- and *Interpretation of the Revised Gesell and Amatruda Developmental Examination*. Hagerstown, MD: Harper & Row; 1980
59. McCabe A, Rollins PR. Assessment of pre-school narrative skills. *Am J Speech Lang Pathol*. 1994;3(1):45–56
 60. Atkinson J, Anker S, Rae S, Hughes C, Braddock O. A test battery of child development for examining functional vision (ABCDEFV). *Strabismus*. 2002;10(4):245–269
 61. Carruth BR, Skinner JD. Feeding behaviors and other motor development in healthy children (2-24 months). *J Am Coll Nutr*. 2002;21(2):88–96
 62. Kimmmerle M, Ferre CL, Kotwica KA, Michel GF. Development of role-differentiated bimanual manipulation during the infant's first year. *Dev Psychobiol*. 2010;52(2):168–180
 63. Egan DF, Brown R. Developmental assessment: 18 months to 4 1/2 years. Performance tests. *Child Care Health Dev*. 1986;12(5):339–349
 64. Williams PG, Lerner MA; Council on Early Childhood; Council on School Health. School readiness. *Pediatrics*. 2019;144(2):e20191766
 65. Kitsao-Wekulo P, Holding P, Abubakar A, et al. Describing normal development in an African setting: The utility of the Kilifi Developmental Inventory among young children at the Kenyan coast. *Learn Individ Differ*. 2016;46:3–10
 66. Gajewska E, Sobieska M, Moczko J, Kuklińska A, Laudńska-Krzemińska I, Osiński W. Independent reaching of the sitting position depends on the motor performance in the 3rd month of life. *Eur Rev Med Pharmacol Sci*. 2015;19(2):201–208
 67. Cox CA, Zinkin PM, Grimsley MF. Aspects of the six-month developmental examination in a longitudinal study. *Dev Med Child Neurol*. 1977;19(2):149–159
 68. World Health Organization Multicentre Growth Reference Study Group. WHO Motor Development Study: windows of achievement for six gross motor development milestones. *Acta Paediatr Suppl*. 2006;95(S450):86–95
 69. Noller K, Ingrisano D. Cross-sectional study of gross and fine motor development. Birth to 6 years of age. *Phys Ther*. 1984;64(3):308–316
 70. American Academy of Pediatrics. *Bright Futures Pocket Guide*, 4th ed. Itasca, IL: American Academy of Pediatrics; 2017
 71. Ohtoshi T, Muraki T, Takada S. Investigation of age-related developmental differences of button ability. *Pediatr Int*. 2008;50(5):687–689
 72. Eremita M, Semancik E, Lerer T, Dworkin PH. Can we identify parents who do not verbally share concerns for their children's development? *J Dev Behav Pediatr*. 2017;38(3):224–227
 73. Sices L, Egbert L, Mercer MB. Sugar-coaters and straight talkers: communicating about developmental delays in primary care. *Pediatrics*. 2009;124(4):e705–e713
 74. Zuckerman KE, Sinche B, Mejia A, Cobian M, Becker T, Nicolaidis C. Latino parents' perspectives on barriers to autism diagnosis. *Acad Pediatr*. 2014;14(3):301–308
 75. Graybill E, Self-Brown S, Lai B, et al. Addressing disparities in parent education: examining the effects of learn the signs/act early parent education materials on parent outcomes. *Early Child Educ J*. 2016;44(1):31–38
 76. Kelly Y, Sacker A, Schoon I, Nazroo J. Ethnic differences in achievement of developmental milestones by 9 months of age: the Millennium Cohort Study. *Dev Med Child Neurol*. 2006;48(10):825–830
 77. Gesell A. *Infancy and Human Growth*. New York, NY: Macmillan; 1928
 78. Lipkin PH, Macias MM, Baer Chen B, et al. Trends in pediatricians' developmental screening: 2002-2016. *Pediatrics*. 2020;145(4):e20190851
 79. Boyle CA, Boulet S, Schieve LA, et al. Trends in the prevalence of developmental disabilities in US children, 1997-2008. *Pediatrics*. 2011;127(6):1034–1042
 80. McManus BM, Richardson Z, Schenkman M, Murphy N, Morrato EH. Timing and intensity of early intervention service use and outcomes among a safety-net population of children. *JAMA Netw Open*. 2019;2(1):e187529
 81. Richardson ZS, Scully EA, Dooling-Litfin JK, et al. Early intervention service intensity and change in children's functional capabilities. *Arch Phys Med Rehabil*. 2020;101(5):815–821
 82. Shonkoff JP, Hauser-Cram P. Early intervention for disabled infants and their families: a quantitative analysis. *Pediatrics*. 1987;80(5):650–658
 83. Guralnick MJ. Effectiveness of early intervention for vulnerable children: a developmental perspective. *Am J Ment Retard*. 1998;102(4):319–345
 84. Rosenberg SA, Zhang D, Robinson CC. Prevalence of developmental delays and participation in early intervention services for young children. *Pediatrics*. 2008;121(6):e1503–e1509
 85. Rosenberg SA, Robinson CC, Shaw EF, Ellison MC. Part C early intervention for infants and toddlers: percentage eligible versus served. *Pediatrics*. 2013;131(1):38–46
 86. Kogan MD, Vladutiu CJ, Schieve LA, et al. The prevalence of parent-reported autism spectrum disorder among US children. *Pediatrics*. 2018;142(6):e20174161
 87. Barger B, Rice C, Wolf R, Roach A. Better together: developmental screening and monitoring best identify children who need early intervention. *Disabil Health J*. 2018;11(3):420–426

Supplemental Information

SUPPLEMENTAL TABLE 7 Deleted Developmental Milestones for the CDC *Learn the Signs. Act Early.* Program

Checklist	Age	Deleted Developmental Milestones	Reasons Removed ^a
1	2 mo	Can briefly calm himself (may bring hands to mouth and suck on hand)	1, 4
2	2 mo	Begins to act bored (cries or fussy) if activity does not change	1, 4
3	2 mo	Makes smoother movements with arms and legs	1
4	4 mo	Copies some movements and facial expressions, like smiling or frowning	1
5	4 mo	Begins to babble	4
6	4 mo	Babbles with expression and copies sounds he hears	1, 4
7	4 mo	Cries in different ways to show hunger, pain, or being tired	1
8	4 mo	Lets you know if she is happy or sad	4
9	4 mo	Responds to affection	4
10	4 mo	Reaches for toy with one hand	3
11	4 mo	Watches faces closely	3
12	4 mo	Follows moving things with eyes from side to side	4
13	4 mo	Recognizes familiar people and things at a distance	3
14	4 mo	Pushes down on legs when feet are on a hard surface	1, 4
15	6 mo	Likes to play with others, especially parents	3
16	6 mo	Responds to other people's emotions and often seems happy	3
17	6 mo	Strings vowels together when babbling (ah, eh, oh) and takes turns with parent while making sounds	3, 4
18	6 mo	Makes sounds to show joy and displeasure	4
19	6 mo	Begins to say consonant sounds (jabbering with m and b)	4
20	6 mo	Begins to pass things from one hand to another	3, 4
21	6 mo	Rolls over in both directions (front to back, back to front)	2
22	6 mo	When standing, supports weight on legs and might bounce	1
23	6 mo	Rocks back and forth, sometimes crawling backward before moving forward	1
24	9 mo	Has favorite toys	1
25	9 mo	Copies sounds and gestures of others	3, 4
26	9 mo	Uses fingers to point at things	3
27	9 mo	Watches the path of something as it falls	1
28	9 mo	Puts things in her mouth	3
29	9 mo	Stands, holding on	2
30	9 mo	Pulls to stand	3
31	9 mo	Crawls	1
32	12 mo	Is shy or nervous with strangers	3
33	12 mo	Has favorite things and people	1
34	12 mo	Shows fear in some situations	3
35	12 mo	Hands you a book when he wants to hear a story	1
36	12 mo	Repeats sounds or actions to get attention	4
37	12 mo	Plays games such as peek-a-boo and pat-a-cake	3
38	12 mo	Makes sounds with changes in tone (sounds more like speech)	1, 4
39	12 mo	Says "mama" and "dada" and exclamations like "uh-oh!"	1
40	12 mo	Tries to say words you say	1, 4
41	12 mo	Explores things in different ways, like shaking, banging, throwing	1
42	12 mo	Finds hidden things easily	4
43	12 mo	Copies gestures	4
44	12 mo	Lets things go without help	1, 4
45	12 mo	Pokes with index (pointer) finger	1, 4
46	12 mo	Follows simple instructions, like "Pick up the toy."	3
47	12 mo	Gets to a sitting position without help	3
48	12 mo	May stand alone	4
49	18 mo	Likes to hand things to others as play	1
50	18 mo	May have temper tantrums	1, 4
51	18 mo	May be afraid of strangers	3, 4
52	18 mo	May cling to caregivers in new situations	3, 4
53	18 mo	Says and shakes head no	2
54	18 mo	Knows what ordinary things are for, for example, telephone, brush, spoon	3
55	18 mo	Shows interest in a doll or stuffed animal by pretending to feed	4
56	18 mo	Points to 1 body part	1
57	18 mo	Pulls toys while walking	1

SUPPLEMENTAL TABLE 7 Continued

Checklist	Age	Deleted Developmental Milestones	Reasons Removed ^a
58	24 mo	Gets excited when with other children	1, 4
59	24 mo	Shows more and more independence	1, 4
60	24 mo	Shows defiant behavior (doing what he has been told not to)	1, 4
61	24 mo	Repeats words overheard in a conversation	1
62	24 mo	Points to things in a book	4
63	24 mo	Finds things even when hidden under 2 or 3 covers	1, 4
64	24 mo	Begins to sort shapes and colors	1, 4
65	24 mo	Completes sentences and rhymes in familiar books	1
66	24 mo	Plays simple make-believe	3, 4
67	24 mo	Builds towers of 4 or more blocks	4
68	24 mo	Might use one hand more than the other	1, 4
69	24 mo	Stands on tiptoe	1
70	24 mo	Walks up and down stairs holding on	2
71	24 mo	Throws ball overhand	2, 4
72	24 mo	Makes or copies straight lines and circles	2
73	3 y	Copies adults and friends	3
74	3 y	Shows affection for friends without prompting	1
75	3 y	Understands the idea of mine and his and hers	1
76	3 y	Shows a wide range of emotions	1, 4
77	3 y	May get upset with major changes in routine	1, 4
78	3 y	Follows instructions with 2 or 3 steps	4
79	3 y	Can name most familiar things	4
80	3 y	Understands words like in, on, and under	1, 4
81	3 y	Names a friend	1
82	3 y	Plays make-believe with dolls, animals, and people	3
83	3 y	Does a puzzle with 3 or 4 pieces	2, 4
84	3 y	Understands what "two" means	1, 4
85	3 y	Builds a tower of more than 6 blocks	1, 4
86	3 y	Climbs well	4
87	3 y	Runs easily	3, 4
88	3 y	Pedals a tricycle (3-wheel bike)	1, 4
89	3 y	Walks up and down stairs, one foot on each step	1, 4
90	4 y	Enjoys doing new things	1
91	4 y	Is more and more creative with make-believe play	4
92	4 y	Cooperates with other children	4
93	4 y	Often cannot tell what is real and what is make-believe	1, 4
94	4 y	Talks about what she likes and what she is interested in	1
95	4 y	Knows some basic rules of grammar, such as correctly using he and she	2, 4
96	4 y	Tells stories	3
97	4 y	Can say first and last name	3
98	4 y	Understands the idea of counting	1, 4
99	4 y	Understands the idea of same and different	1, 4
100	4 y	Uses scissors	2, 4
101	4 y	Starts to copy some capital letters	4
102	4 y	Plays board or card games	1
103	4 y	Tells you what he thinks is going to happen next in a book	4
104	4 y	Pours, cuts with supervision, and mashes own food	1
105	5 y	Wants to please friends	4
106	5 y	Wants to be like friends	4
107	5 y	Is aware of gender	4
108	5 y	Can tell what is real and what is make-believe	1, 4
109	5 y	Shows more independence (eg, may visit a next-door neighbor by himself [adult supervision is still needed])	1, 4
110	5 y	Is sometimes demanding and sometimes very cooperative	1, 4
111	5 y	Speaks very clearly	3
112	5 y	Uses future tense; for example, "Grandma will be here."	1
113	5 y	Says name and address	1
114	5 y	Can draw a person with at least 6 body parts	2
115	5 y	Copies a triangle and other geometric shapes	2, 4
116	5 y	Knows about things used every day, like money and food	1, 4
117	5 y	Stands on one foot for 10 s or longer	1, 4

SUPPLEMENTAL TABLE 7 Continued

Checklist Age		Deleted Developmental Milestones	Reasons Removed ^a
118	5 y	Hops; may be able to skip	1, 4
119	5 y	Can do a somersault	1, 4
120	5 y	Uses a fork and spoon and sometimes a table knife	1, 4
121	5 y	Can use the toilet on her own	2, 4
122	5 y	Swings and climbs	1, 4

^a Key: 1 = little/no normative data regarding when the milestone/part of the milestone should be achieved by $\geq 75\%$ of children; 2 = inconsistent data regarding when the milestone/part of the milestone should be achieved by $\geq 75\%$ of children; 3 = duplicate milestone at earlier or later age; 4 = did not meet other criteria established by SMEs (see Table 1 for criteria).